

## Mortality due to COVID-19 during the vaccination plan against the SARS-CoV-2 virus in Cali, Colombia

### Mortalidad por COVID-19 durante el plan de vacunación contra el virus SARS-CoV2 en Cali, Colombia

Carlos Reina,<sup>1</sup>  Miyerlandi Torres,<sup>1</sup>  Edgar Muñoz,<sup>2</sup>  Elvia Karina Grillo Ardila,<sup>3</sup>  Rodrigo Guerrero Velasco,<sup>4</sup>  Luis Eduardo Bravo<sup>5,6</sup>   
[carlos.reina@udea.edu.co](mailto:carlos.reina@udea.edu.co)

**1** Secretaria de Salud Pública del Municipio de Santiago de Cali, Colombia, **2** University of Texas , Health Science Center San Antonio, Texas, USA, **3** Universidad del Valle, Facultad de Salud, Doctorado en Salud, Cali, Colombia., **4** Universidad del Valle, Instituto de Investigación y Desarrollo en Prevención de la Violencia y Promoción de la Convivencia Social, CISALVA, Cali, Colombia., **5** Universidad del Valle, Facultad de Salud, Escuela de Medicina, Departamento de Patología, Cali, Colombia., **6** Hospital Universitario del Valle “Evaristo García”, Cali, Colombia.



#### OPEN ACCESS

**Citation:** Reina C, Torres M, Muñoz E, Grillo AEK, Guerrero VR, Bravo LE. Mortality due to COVID-19 during the vaccination plan against the SARS-CoV-2 virus in Cali, Colombia. Colomb Méd (Cali), 2021; 52(4):e7005047 <http://doi.org/10.25100/cm.v52i4.5047>

#### Keywords:

COVID-19; mortalidad; SARS-CoV-2, mortalidad; vacunas; vacunación.

#### Palabras clave:

Covid-19; mortalidad; SARS-CoV-2, mortalidad; vacunas; vacunación.

**Copyright:** © 2021 Universidad del Valle



#### Conflicts of interest:

The authors declare no conflicts of interest

#### Corresponding author:

Carlos Reina. E-mail: [carlos.reina@udea.edu.co](mailto:carlos.reina@udea.edu.co)


Dear Editor:

The behavior of the COVID-19 pandemic during the three peaks registered in Cali has been heterogeneous, and the severity of infections and deaths has varied over time (Figure 1). Cali, with 2.2 million inhabitants, is the third most populated city in Colombia <sup>1</sup>; as of November 21, 2021, it has recorded 285 thousand confirmed cases and 7,474 deaths due to the infection <sup>2</sup>. The vaccination plan against SARS-CoV-2, which was started in February, has immunized 1.54 million people in Cali <sup>3</sup>.

As in the rest of the country, the development of the vaccination plan in Cali coincided with the third peak of the pandemic, the most lethal and prolonged; as well as with multiple social mobilizations of hundreds of protesters who crowded the streets, between May and June 2021, protesting against an unpopular tax reform enacted by the national government. In Cali, the vaccination plan was also affected by lack of access to health services and vaccination points due to blockages registered in different zones of the city. At the end of the third peak, the country faced difficulties in continuing with the vaccination plan due to barriers to accessing vaccines in the international market.

In accordance with the national mandate, all SARS-CoV-2 virus infections diagnosed in the country's private and state laboratories must be notified to the Public Health Secretaries. In case of doubt, the territorial entity, the municipality of Cali in our case, analyzed each of the deaths and determined which ones were caused by COVID-19. From February to November 2021, city laboratories reported 139,972 SARS-CoV-2 infections and 3,312 deaths for COVID-19. This figure corresponds to 64% of the deaths that occurred in Cali during the pandemic. 82.5% of infections and 89.1% of deaths occurred in persons with incomplete vaccination schemes or unvaccinated people.

As it has been reported in other countries, the effectiveness of vaccines to prevent deaths from COVID-19 is variable and lower than that observed in clinical trials <sup>4</sup>. The effectiveness of vaccines against SARS-CoV-2 was estimated with the hazard ratio (HR) using a Cox proportional hazards model considering as a predictor covariate the vaccination status of each



**Figure 1.** Cali, Colombia. Mortality due to COVID-19 during the development of the national vaccination plan between April 2020 and November 2021. Monthly moving average. The first peak began on June 7 and ended on August 13, 2020. The second peak began on December 13, 2020; and ended on February 3, 2021. The third peak began on March 25 and ended on August 8, 2021; it was the deadliest and caused half of the deaths of the entire pandemic in Cali.

individual in different times. Changes in HR associated with partial immunization ( $\geq 14$  days after receiving the first dose and before receiving the second dose) and complete immunization ( $\geq 14$  days after receiving the second dose), further adjusting for age, sex, presence of comorbidities and vaccination week.

Mortality from COVID-19 in the city of Cali changed / dropped from a daily average of 15 deaths in November 2020 to 3 in November 2021. The risk of dying was higher in the unvaccinated, in men compared with women (HR: 1.96, CI 95%: 1.83-2.10), and in people with comorbidities (HR: 2.74, 95% CI: 2.55-2.94). An 8% increase for each year of age was evident. Of special interest was to observe variations according to the type of vaccine Table 1.

Adjusting for vaccination status, age, sex, presence of comorbidities and vaccination week. CI: confidence interval.

**Table 1.** Cali, Colombia. Effectiveness of vaccines against SARS-CoV-2

Variable		HR	CI(95%)	Effectiveness %	CI (95%)
Vaccine	Not vaccinated	1			
	AstraZeneca®	0.40	(0.30-0.53)	59.7	(48.3-71.1)
	Coronavac®	0.38	(0.29-0.49)	62.4	(52.6-72.3)
	Janssen®	0.11	(0.05-0.25)	89.1	(80.2-98.0)
	Moderna®	0.02	(0.01-0.09)	97.8	(94.8-100.8)
	Pfizer®	0.26	(0.19-0.35)	74.4	(66.3-82.6)
Vaccination status	Not vaccinated	1			
	Complete immunization	0.12	(0.07-0.20)		
Vaccination week	Semanas desde inicio del plan	0.99	(0.98-0.99)		
Sexo	Female	1			
	Male	1.96	(1.83-2.10)		
Age	Years	1.08	(1.07-1.08)		
Comorbidity	Absent	1			
	Present	2.74	2.55-2.94		

Adjusting for vaccination status, age, sex, presence of comorbidities and vaccination week. CI: confidence interval. Effectiveness of each vaccine (EV) = 1 - HR, where HR is the exponential of the coefficient associated with the vaccine in the previously adjusted model. The confidence intervals of EV were estimated using the delta method<sup>5</sup>.

Effectiveness of each vaccine (EV) =  $1 - HR$ , where HR is the coefficient associated with the vaccine in the previously adjusted model. The confidence intervals of EV were estimated using the delta method <sup>5</sup>

Although vaccines against the SARS-CoV-2 virus showed 100% efficacy to prevent death, and 85-90% for severe disease in clinical trials <sup>6</sup>, the effectiveness reported in the real world has been lower <sup>7</sup>. Table 1 shows that in Cali, the effectiveness to prevent death from COVID-19 was lower than that reported in clinical trials; in addition, it showed variations according to the type of vaccine that was administered: 60% for *AstraZeneca*, 62% for *CoronaVac*, 89% for *Janssen*, 98% for *Moderna*, and 74% for *Pfizer*. Part of this variability may be due to the availability of vaccines and failures in compliance with the established times to adequately complete the vaccination schedule, as well as the capacities to develop the vaccination plan.

The conditions described in Cali and Colombia during the third peak could have favored the appearance of more contagious variants of the SARS-CoV-2 virus, but which could be controlled by current vaccines, probably because they evolved in an unvaccinated population <sup>8</sup>. The *a priori* reasoning is that those who have been vaccinated and/or have recovered favor the appearance of **collective** herd immunity. The ones who die increase the percentage of vaccinated people, by reducing the unvaccinated population. However, the equation is more complex because unvaccinated groups can be reservoirs for the appearance of new variants that have evolved in the vaccinated population, and that be resistant to current vaccines, making those already vaccinated susceptible again.

In this globalized and interconnected world, herd immunity cannot be considered for isolated populations. There are differences in the variants between countries. While in Colombia, as of November 2021, the DELTA variant was the dominant one <sup>9</sup>; in South Africa, a country with low vaccination coverage, it has emerged theOMICRON variant with 26 new mutations <sup>10</sup>. It is urgent to accelerate the rate of vaccination to stop the chain of transmission and reduce the probability of the appearance of new variants. Further studies are required to determine in other regions the effectiveness of the various vaccines used in Colombia, considering the level of adherence and access to them.

## References

1. Departamento Administrativo Nacional de Estadística (DANE). Proyecciones de población 2018-2023. DANE; 2020. Cited: 2021 Nov 21. Available from: <https://www.dane.gov.co/index.php/estadisticas-por-tema/demografia-y-poblacion/proyecciones-de-poblacion>.
2. Secretaria de Salud Pública. COVID19\_21noviembre. Boletín Epidemiológico No.622. Cali: Alcaldía de Santiago de Cali; 2021. Available from: <https://www.cali.gov.co/salud/publicaciones/152840/boletines-epidemiologicos/>.
3. Secretaria de Salud Pública. COVID19\_23noviembre. Boletín Vacunación COVID No.265. Cali: Alcaldía de Santiago de Cali; 2021. Cited: 2021 Nov 21. Available from: <https://www.cali.gov.co/salud/publicaciones/159073/boletines-vacunacion-covid/>
4. Jara A, Undurraga EA, González C, Paredes F, Fontecilla T, Jara G, et al. Effectiveness of an Inactivated SARS-CoV-2 Vaccine in Chile. *N Engl J Med*. 2021; 385: 875-884. DOI: 10.1056/NEJMoa2107715. [PubMed]
5. Thompson MG, Burgess JL, Naleway AL, Tyner HL, Yoon SK, Meece J, et al. Interim estimates of vaccine effectiveness of BNT162b2 and mRNA-1273 COVID-19 vaccines in preventing SARS-CoV-2 infection among health care personnel, first responders, and other essential and frontline workers - eight u.s. locations, December 2020-March . *MMWR Morb Mortal Wkly Rep*. 2021;70(13):495-500. [PubMed]

6. Kai X, Xiao-Yan T, Miao L, Zhang-Wu L, Jiang-Nan C, Jiao-Jiao L, et al. Efficacy and safety of COVID-19 vaccines: A systematic review. *Chinese J Contemp Pediatr*. 2021; 23(3):221-8. doi: 10.7499/j.issn.1008-8830.2101133
7. Higdon MM, Wahl B, Jones CB, Rosen JG, Truelove SA, Baidya A, et al. A systematic review of COVID-19 vaccine efficacy and effectiveness against SARS-CoV-2 infection and disease. *medRxiv*. 2021. Doi: 10.1101/2021.09.17.21263549
8. Goldman E. How the unvaccinated threaten the vaccinated for COVID-19: A Darwinian perspective. *Proc Natl Acad Sci United States Am*. 2021;118(39): e2114279118. Doi: 10.1073/pnas.2114279118.
9. Instituto Nacional de Salud. COVID-19 en Colombia; 2021. Cited: 2021 Dec 14. Available from: <https://www.ins.gov.co/Noticias/Paginas/coronavirus-genoma.aspx>
10. Mullen JL, Tsueng G, Latif AA, Alkuzweny M, Cano M, Haag E, et al. Omicron variant report. *outbreak.info, Lineage | Mutation Tracker*; 2021. Cited: 2021 Dec 14. Available from: <https://outbreak.info/situation-reports/omicron>